

2-STACK SORTABLE PERMUTATIONS WITH A GIVEN NUMBER OF RUNS

MIKLÓS BÓNA

ABSTRACT. Using earlier results we prove a formula for the number $W_{(n,k)}$ of 2-stack sortable permutations of length n with k runs, or in other words, $k - 1$ descents. This formula will yield the suprising fact that there are as many 2-stack sortable permutations with $k - 1$ descents as with $k - 1$ ascents. We also prove that $W_{(n,k)}$ is unimodal in k , for any fixed n .

1. INTRODUCTION

1.1. Our main results. In this paper we are going to show that the number of 2-stack sortable permutations of length n with $k - 1$ ascents is equal to the number $T_{(n,k)}$ of $\beta(1, 0)$ -trees on $n + 1$ nodes with k leaves. (See Section 2 for the definition of $\beta(1, 0)$ -trees). This, and results from [2] and [5] will enable us to easily show that

$$W_{(n,k)} = \frac{(n + k - 1)!(2n - k)!}{k!(n + 1 - k)!(2k - 1)!(2n - 2k + 1)!}, \quad (1)$$

which formula is symmetric in k and $n + 1 - k$.

1.2. Background and Definitions. In what follows, permutations of length n will be called n -permutations. We say that i is a *descent* of the permutation $p = (p_1, p_2, \dots, p_n)$ if $p_i > p_{i+1}$. Similarly, j is an *ascent* of p if $p_j < p_{j+1}$. If p has d descents, then p decomposes into $d + 1$ increasing sequences of consecutive entries, and we then say that p has $d + 1$ *runs*. We will use the concept of descents and runs interchangeably, according to the current context.

The stack-sorting operation Π can be defined on the set of all n -permutations as follows. Let $p = p_L n p_R$ be an n -permutation, with p_L and p_R respectively denoting its subword before and after the maximal entry. Let $\Pi(p) = \Pi(p_L)\Pi(p_R)n$, where p_L and p_R are defined recursively by this same rule. For a nonrecursive, algorithmic definition, or the origin of the notion see [6], [8].

A permutation p is called t -stack sortable if $\Pi^t(p)$ is the identity permutation.

The set of 1-stack sortable n -permutations is easy to characterize by the following notion of pattern avoidance. Let $q = (q_1, q_2, \dots, q_k)$ be a k -permutation and let $p = (p_1, p_2, \dots, p_n)$ be an n -permutation. We say that p contains a q -subsequence (or q -pattern) if there exists $1 \leq i_{q_1} < i_{q_2} < \dots < i_{q_k} \leq n$ such that $p_{i_1} < p_{i_2} < \dots < p_{i_k}$. We say that p avoids q if p contains no q -subsequence.

This paper was written while the author was a one-term visitor at Mathematical Sciences Research Institute in Spring 1997. This visit was supported by an MIT Applied Mathematics Fellowship.

For example, p avoids 231 if it cannot be written as $\cdots, a, \cdots, b, \cdots, c, \cdots$ so that $c < a < b$. It is easy to show [6] [8] [9] that a permutation is 1-stack sortable if and only if it avoids the pattern 231. In particular, the number of 1-stack sortable permutations is therefore C_n , the n -th Catalan number.

The set of 2-stack sortable permutations is much more complex. For example, it is not true that a subword of a 2-stack sortable permutation is always 2-stack sortable; for 35241 is 2-stack sortable, while 3241 is not. Similarly, it is far more difficult to find a formula for the number W_n of these permutations than for that of 1-stack sortable ones. See [10] [4] or [3] for a proof of the formula $W_n = 2(3n)!/((n+1)!(2n+1)!)$ and see [8] for a characterization of 2-stack sortable permutations.

In this paper we are going to construct a bijection between the set of 2-stack sortable n -permutations with k runs and that of $\beta(1, 0)$ -trees on $n+1$ vertices having k leaves. This bijection will be based on a bijection in [4], which mapped these permutations into a class of labeled binary trees. (Our trees are not binary). The set of these trees has recently been shown to be equinumerous [2] to a certain class of planar maps, and the number of those planar maps is known [5]. Therefore, we are going to obtain a *formula* for the number $W_{(n,k)}$ of 2-stack sortable permutations with k runs, which is a bit surprising, given that even the formula for W_n was fairly difficult to prove.

What is even more surprising is that this formula reveals that *there are as many 2-stack sortable permutations with k descents as with k ascents*. We do not see any obvious reason why this should be true. A direct combinatorial proof of this fact would be highly desirable, but it may not be easy as we do not even have such a proof for the same fact in the much simpler case of 132-avoiding permutations. We also prove that $W_{(n,k)}$ is unimodal in k for any fixed n .

We point out that this is the second case recently [1] when a class of the description trees introduced in [2] to enumerate planar maps has been used to enumerate permutations.

2. THE CORRESPONDENCE BETWEEN TREES AND PERMUTATIONS

Let p be an n -permutation. A *right-to-left maximum* of p is an entry which is larger than all entries it precedes. If the right-to-left maxima of p are $a_1 > a_2 > \cdots > a_t$, then clearly $a_1 = n$ and a_t is the last entry of p . In the next three paragraphs we closely follow the characterization of 2-stack sortable permutation given in [4].

Write p in the form $s_1 a_1 s_2 a_2 \cdots s_t a_t$. Here the s_i are strings of entries located between two right-to-left maxima. We say that p is of type 1 if the entry $a_t - 1$ is part of the string s_t . Otherwise we say that p is of type 2. The 1-permutation 1, denoted ϵ , is of type 2. Then the following holds.

Lemma 1. [4] *There is a bijection F from the set of 2-stack sortable n -permutations of type 1 to that of 2-stack sortable $(n-1)$ -permutations with a right-to-left maximum marked. Moreover, F preserves the number of descents and does not decrease the number of right-to-left maxima.*

Proof:[4] Let p be of type 1, with the above $s_1 a_1 s_2 a_2 \cdots s_t a_t$ decomposition. Define F as follows. Delete a_t from the end of p and decrement all entries larger than a_t

by 1. Then $a_t - 1 \in s_t$ becomes a right-to-left maximum in the new permutation, in fact, it becomes the t th right-to-left maximum. Mark this entry to get $F(p)$. This map is reversible by simply adding an entry one larger than the marked vertex to the end of $F(p)$ and increment all the larger entries by one.

So F is a bijection. Moreover, F does not create or destroy any existing descents as it deletes the last entry of p , and that entry was larger than its predecessor. The number of right-to-left maxima is not decreased by F , because $a_t - 1$ becomes a right-to-left maximum instead of a_t .

Finally, one sees easily [4] that $F(p)$ is 2-stack sortable if and only if p is, and the proof is complete. $\square$

We need one more lemma from [4]. We omit its proof as we will not need any of its rather complicated machinery. Let $rl(p)$ be the number of right-to-left maxima of p and let $desc(p)$ be the number of descents in p .

Lemma 2. *Let $n \geq 2$. Then there is a bijection G from the set of 2-stack sortable n -permutations of type 2 onto the set of ordered pairs (p_1, p_2) where*

1. p_1 is any 2-stack sortable n_1 -permutation, p_2 is either a 2-stack sortable n_2 -permutation of type 1, or ϵ ,
2. $n_1 + n_2 = n$,
3. $rl((p_1) + rl(p_2) = rl(p)$,
4. $desc(p_1) + desc(p_2) + 1 = desc(p)$.

We are going to iterate the decomposition of the above Lemma. In other words, we take a 2-stack sortable n -permutation p of type 2, decompose it into (p_1, p_2) by the bijection G of [4], then, if p_1 was of type 2 (and not ϵ), then we apply G to p_1 to get (p_{11}, p_{12}) . Again, if p_{11} is of type 2 (and not ϵ), then we apply G to p_{11} , and so on. We stop when both elements of the current decomposition are either of type 1, or ϵ . As each step of this algorithm uses a bijection, p can be recovered from its final decomposition. This proves the following Corollary.

Corollary 1. *There is a bijection H from the set of 2-stack sortable n -permutations onto the set of string $(q_1, q_2, \dots, q_r)$, where $r \geq 1$ and*

- for all i , q_i is either a 2-stack sortable permutation of type 1, or ϵ and
- $\sum_{i=1}^r |q_i| = n$,
- $\sum_{i=1}^r rl(q_i) = rl(p)$,
- the total number of runs in the q_i equals the number of runs in p .

The last part follows from the last part of Lemma 2 and the fact that the number of runs in any permutation is one more than that of descents.


Cori, Jacquard and Schaeffer [2] give the following definition in their study of planar maps.

Definition 1. *A rooted plane tree with nonnegative integer labels $l(v)$ on each of its vertices v is called a $\beta(1, 0)$ -tree if it satisfies the following conditions:*

- if v is a leaf, then $l(v) = 1$,
- if v is the root and $v_1, v_2, \dots, v_k$ are its children, then $l(v) = \sum_{i=1}^k l(v_i)$,

- if v is a nonroot internal node and $v_1, v_2, \dots, v_k$ are its children, then $l(v) \leq \sum_{i=1}^k l(v_i)$.

Example 1. Two $\beta(1, 0)$ -trees are shown in Figures 1a and 1b.



Now we are in position to state and prove the main result of this paper.

Theorem 1. *For any positive integer n and for any positive integer $k \leq n$, there is a bijection B from the set of 2-stack sortable n -permutations with k runs onto that of $\beta(1, 0)$ -trees on $n + 1$ nodes having k leaves. Therefore, $W_{(n,k)} = T_{(n,k)}$. Moreover, for any 2-stack sortable n -permutation p , the label of the root of $B(p)$ equals $rl(p)$.*

Proof: We are going to prove the statements of the Theorem by induction on n and k . For $k = 1$ and any n , the statements are clearly true: there is one n -permutation with one run, namely $12 \cdots n$, and there is one $\beta(1, 0)$ -tree having only one leaf, namely the one consisting of a single path. The root label of this tree is 1, and the number of right-to-left maxima of this permutation is 1, so the initial case is proved.

Now let $k > 1$ and suppose we already know the statements for all permutations shorter than n . Let p be any 2-stack sortable n -permutation with k runs.

- First suppose p is of type 1. Consider $F(p)$, where F is the bijection defined in the proof of Lemma 1. Forget for now which vertex of $F(p)$ is marked, then $F(p)$ is a 2-stack sortable $(n - 1)$ -permutation with k runs. So, by our induction hypothesis, B associates a $\beta(1, 0)$ -tree T' on n nodes to it, which has k leaves and whose root label is equal to $rl(F(p)) \geq rl(p)$. Now add one node above the root x of T' and “recall” which right-to-left maximum of $F(p)$ was marked. If it was the t th right-to-left maximum, that is, if p had exactly t right-to-left maxima, then let the label of this new root x (and necessarily, that of the old root, which is the only child of x) be t . Then we certainly get a $\beta(1, 0)$ -tree T as $t \leq rl(F(p))$. We set $B(p) = T$. By the argument of Lemma 1 we see that p can be recovered from T (as T' , and thus $F(p)$ can), and that T has k leaves, and its root label is $rl(p)$ as it should be.
- If $p = \epsilon$, then $B(p)$ is the only $\beta(1, 0)$ -tree on two nodes. We point out that here, as well as in the previous case, the root of $B(p)$ has only one child.
- Now suppose p is of type 2. Take $H(p) = (q_1, q_2, \dots, q_r)$ as defined in Corollary 1 and consider $B(q_i)$ for all i as defined by the previous two cases. $B(q_i)$ has $q_i + 1$ nodes, so by contracting the roots of all $B(q_i)$ to one node (keeping the

$B(q_i)$ ordered from left to right), we get a tree T with $1 + \sum_{i=1}^r |q_i| = n + 1$ nodes. We set $B(p) = T$. The components q_i can be recovered from T by simply cutting off its root and adding a new root to the top of each component obtained. (Recall that in all the $B(q_i)$, the root has only one child). The number of leaves of T equals the sum of the leaves of the $B(q_i)$, so by the previous two cases and by Corollary 1, it equals the number of runs of p . The root label of T is by definition the sum of the labels of its children, which is, again by the previous two cases and by Corollary 1, equal to $rl(p)$.

This completes the proof of the Theorem. $\square$

We remark that our algorithm certainly sets up a bijection between the set of *all* 2-stack sortable n -permutations and that of *all* $\beta(1, 0)$ -trees on $n + 1$ vertices.

3. NEW ENUMERATIVE RESULTS

Exercise 2.9.8.b in [5] shows that the number of nonseparable rooted planar maps with $f + 1$ faces and $p + 1$ vertices is equal to

$$\frac{(2f + p - 2)!(2p + f - 2)!}{f!p!(2f - 1)!(2p - 1)!}. \quad (2)$$

It is shown in [2] that the number of these maps equals the number of $\beta(1, 0)$ -trees on $n + 1$ nodes with $k = f + 1$ leaves and $n + 1 - k = p + 1$ internal nodes. (Here the root is counted as an internal node). By Theorem 1 this implies the result we announced in the introduction.

Theorem 2. *Let $W_{(n,k)}$ be the number of 2-stack sortable n -permutations with k runs. Then for all $1 \leq k \leq n$ we have*

$$W_{(n,k)} = \frac{(n + k - 1)!(2n - k)!}{k!(n + 1 - k)!(2k - 1)!(2n - 2k + 1)!}.$$

In particular, equation (2) shows that the roles of $f + 1 = k$ and $p + 1 = n + 1 - k$ are symmetric. In other words, $W_{(n,k)} = W_{(n,n+1-k)}$. A permutation with k runs has $k - 1$ descents, so have proved the following surprising Corollary.

Corollary 2. *The number of 2-stack sortable n -permutations with k descents equals that of those with k ascents.*

A routine computation yields that $W_{(n,k)}/W_{(n,k-1)} > 1$ if and only if $k \leq (n + 1)/2$, and we already know from the previous Corollary that if $n = 2k$, then $W_{(n,k)} = W_{(n,k+1)}$. So we have proved the following Corollary.

Corollary 3. *For fixed n , the sequence $W_{(n,k)}$ is unimodal in k , and its peak is at $k = \lfloor (n + 1)/2 \rfloor$.*

ACKNOWLEDGEMENT

I am grateful to Gilles Schaeffer who sent me the preprint [2].

REFERENCES

- [1] M. Bóna, Exact enumeration of 1342-avoiding permutations; A close link with labeled trees and planar maps, *Journal of Combinatorial Theory, Series A*, to appear.
- [2] R. Cori, B. Jacquard, G. Schaeffer, Description trees for some families of planar maps, Proceedings of the 9th Conference of Formal Power Series and Algebraic Combinatorics, to appear.
- [3] S. Dulucq. S. Gire, J. West, Permutations with forbidden subsequences and nonseparable planar maps. Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993), *Discrete Math.* **153** (1996), 85-103.
- [4] I. P. Goulden, J. West, Raney paths and a combinatorial relationship between rooted nonseparable planar maps and two-stack-sortable permutations, *J. Combin. Theory Ser. A* **75** (1996), 220-242.
- [5] I. P. Goulden, D. M. Jackson “Combinatorial Enumeration”, Wiley-Interscience, 1983.
- [6] D. E. Knuth, “The art of computer programming,” volume 3, Sorting and searching, Addison-Wesley, Reading, Massachusetts, 1973.
- [7] J. W. Tutte, A census of planar maps, *Canadian Journal of Mathematics* **33** (1963), 249-271.
- [8] J. West, Permutations with forbidden subsequences; and, Stack sortable permutations, PHD-thesis, Massachusetts Institute of Technology, 1990.
- [9] Sorting twice through a stack. Conference on Formal Power Series and Algebraic Combinatorics (Bordeaux, 1991). *Theoret. Comput. Sci.* **117** (1993) 303-313.
- [10] D. Zeilberger, A proof of Julian West’s conjecture that the number of two-stack-sortable permutations of length n is $2(3n)!/((n+1)!(2n+1)!)$. *Discrete Math.* **102** (1992), 85-93.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139